



## **Comparison of Mycelial Growth of Different *Tricholoma matsutake* Strains in Soil Medium at Varying Temperatures**

**Maki Narimatsu<sup>1\*</sup>, Muneyoshi Yamaguchi<sup>2</sup>, Takashi Yamanaka<sup>2</sup>, Seiki Gisusi<sup>3</sup>,  
Tomonori Azuma<sup>3</sup>, Yutaka Tamai<sup>4</sup>, Tohru Fujita<sup>5</sup> and Masataka Kawai<sup>6</sup>**

<sup>1</sup>Iwate Prefectural Forest Technology Center, 560-11 Kemuyama, Yahaba, Iwate 028-3623, Japan.

<sup>2</sup>Forest Research and Management Organization, 1 Matsunosato, Tsukuba, Ibaraki, 305-8687,  
Japan.

<sup>3</sup>Forest Products Research Institute, Hokkaido Research Organization, Nishikagura 1-10, Asahikawa,  
Hokkaido, 071-0198, Japan.

<sup>4</sup>Hokkaido University, Kita 9, Nishi 9, Kita-ku, Sapporo, Hokkaido, 060-8589, Japan.

<sup>5</sup>Kyoto Prefectural Forest Technology Center, Tsuchiya 1, Honjou, Kyoutanba, Funai, Kyoto,  
622-0200, Japan.

<sup>6</sup>Nara Forest Research Institute, Kibi 1, Takatori, Takaichi, Nara, 635-0133, Japan.

### **Authors' contributions**

This work was carried out in collaboration among all authors. Authors MN, MY and TY designed the study. Authors MN and MY carried out the experiment, performed the statistical analysis and wrote the protocol. Authors MN, SG, YT, TF and MK prepared strains for this study. All authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/AJB2T/2019/v5i430066

#### Editor(s):

(1) Dr. Zafar S. Khan, Department of Botany, Maharashtra College of Arts, Science and Commerce, 246-A Jehangir Boman  
Behram Road, Mumbai, India.

#### Reviewers:

(1) Ekachai Chukeatirote, Mae Fah Luang University, Thailand.

(2) Kaushal Kishore Choudhary, Ram Jaipal College, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/52395>

**Short Research Article**

**Received 05 September 2019**

**Accepted 10 November 2019**

**Published 20 November 2019**

### **ABSTRACT**

**Aims:** To investigate the effect of temperature on the mycelial growth of the highly prized mushroom, *Tricholoma matsutake*, in soil medium.

**Methodology:** Seven strains of *T. matsutake* collected from different areas of Japan were incubated in soil medium at 5, 10, 15, 20, 25, and 30°C for 89 day. After incubation, the linear

\*Corresponding author: E-mail: [m-narimatsu@pref.iwate.jp](mailto:m-narimatsu@pref.iwate.jp);

growth of mycelia from the inoculum was measured on vertical lines under a dissecting microscope and classified by the cultivation day number to obtain the growth rate in each vial. The mycelial densities in the soil medium in all the vials were analyzed using qPCR to determine the mycelial biomass.

**Results:** The mycelial growth of *T. matsutake* strains was mainly observed at 5–25°C. The optimum temperatures for linear mycelial growth and for the increase in mycelial density were 19.6°C and 17.6°C, respectively. These values were lower than those reported previously for *T. matsutake* grown on agar medium or in liquid medium. These optimum temperatures showed no clear relationship with the mean air temperature at the site of strain collection.

**Conclusion:** It can be suggested that a temperature around 18°C is suitable for cultivation of *T. matsutake* mycelia in the soil medium.

**Keywords:** *Tricholoma matsutake*; linear mycelial growth; mycelial density; optimum temperature.

## 1. INTRODUCTION

*Tricholoma matsutake* is an ectomycorrhizal fungus in Agaricales, Basidiomycota, with the basidiomata of pileus being ~8–20 cm across and stipe length being ~10–20 cm [1]. *T. matsutake* is an economically important fungus and usually inhabits forests primarily composed of *Pinus densiflora* (Japanese red pine) in Japan [2]. It also grows in the pine forests in other countries [3,4], and substantial amounts of it are exported to Japan [5]. A stable artificial cultivation method for *T. matsutake* has not yet been developed [6]. The fruiting bodies of *T. matsutake* need to be collected from natural forests, which makes its production unsustainable because of the effects of environmental conditions that change annually [7]. For sustainable production of *T. matsutake* fruiting bodies, expanding the area of occurrence of this fungus by inoculation of mycelium or mycorrhizal nursery stock in forests using cultured mycelium is, therefore, essential [5].

In basidiomycetes, the mycelial growth rate, with regard to both the length and density, is affected by temperature [8]. Therefore, temperature is an important factor in determining the culture conditions for basidiomycetes. The effect of temperature on mycelial growth has been reported for saprophytic fungi as well as for ectomycorrhizal fungi [9], including *T. matsutake*. For naturally growing *T. matsutake*, the expansion rate of the fairy ring formed by the fruiting bodies showed a positive relation with the air temperature [7]. This phenomenon suggests that air temperature affects the growth rate of underground mycelia by modulating the soil temperature. Notably, mycelial growth of *T. matsutake* is slower than that of some other ectomycorrhizal fungi, such as *Rhizopogon roseolus* and *Suillus bovinus* [10]. This slower mycelial growth is an obstacle for efficient

experimentation involving *T. matsutake* cultivation.

The optimum temperature for mycelial colony expansion of *T. matsutake* on agar media was reported to be 23°C [11]. However, in nature, *T. matsutake* mycelia grow in the soil [12], and the effect of temperature on mycelial growth in soil remains poorly understood. Furthermore, the optimum temperature for mycelial growth of several saprophytic fungi is known to differ not only among species [13] but also among strains [14]. However, a comparison of the mycelial growth of different *T. matsutake* strains, isolated from various areas of Japan, in the soil medium has not been reported. Therefore, in this study, we measured the mycelial growth, in terms of the linear growth and biomass increase, of different *T. matsutake* strains in the soil at different temperatures.

## 2. MATERIALS AND METHODS

### 2.1 Fungal Strains

Seven *T. matsutake* strains isolated from different areas of Japan were used in this study (Table 1). Among these, two strains were isolated from the Hokkaido Island, three from the northern part of the mainland (e.g., Iwate prefecture), and two from the southern part of the mainland (e.g., Kyoto prefecture and Nara prefecture). The geographic distance between the most distant sites is approximately 1238 km. Living cultures of all the samples have been deposited in the institutions of the authors.

### 2.2 Cultivation of Mycelia

#### 2.2.1 Isolation of mycelia

The strains used in this study were isolated from hymenia except strain NF2970 from lammella,

using nutrient agar media, for example the modified MYPG medium [15] (2.5 g malt extract, 1.0 g yeast extract, 1.0 g peptone, 5.0 g glucose, 15 g agar, and 1000 mL deionized water), and stored at 4°C until use.

### 2.2.2 Inoculation of mycelia

Mycelial growth rate of *T. matsutake* was determined using flat-bottom incubation vials (30 mm outer diameter and 120 mm height) containing the soil medium (Fig. 1) prepared as follows: loamy soil (Turf soil, Koujiya Co. Ltd., Japan), with a particle diameter of 1.4 mm, was soaked in modified MYPG liquid medium [15] (0.25% malt extract, 0.1% yeast extract, 0.1% peptone, 0.5% glucose, pH 5.0) for 16 h to allow the soil to absorb the MYPG medium. Consequently, the soil was set onto a plastic mesh basket for 30 min to remove the extra medium. The B-horizon soil (brown forest soil)


was collected from a depth of 10–20 cm at the *T. matsutake* growth site in the Japanese red pine forest [7], air dried, and passed through a sieve (7-mm mesh size) to remove any stones from the soil. Thereafter, the B-horizon soil was mixed with loamy soil at a ratio of 3:1 (v/v). The moisture content was adjusted to 25% (w/w) with tap water. This soil medium was then filled into the incubation vials to a height of 8 cm (48.5 g) and autoclaved at 121°C for 60 min.

Each strain was pre-incubated on modified Norkrans' C agar medium [16] (1.0 g KH<sub>2</sub>PO<sub>4</sub>, 0.5 g MgSO<sub>4</sub>·7H<sub>2</sub>O, 0.5 mL of 0.2% ZnSO<sub>4</sub>, 0.5 g NH<sub>4</sub>-tartrate, 0.5 mL of 1.0% Fe-citrate, 50 µg thiamine, 0.23 g casein hydrolysate, 0.5 g yeast extract, 10 g glucose, 15 g agar, and 1000 mL deionized water; pH 5.0) at 23°C. After 57 d of incubation, the growth zone of the colony was bored with a cork borer (9-mm diameter), and five mycelial plugs were placed in 125-mL

**Table 1. Study sites and optimum temperature for linear mycelial growth and mycelial density increase**

Strain	Site	Latitude	Longitude	Mean air temperature <sup>(a)</sup>	Optimum temperature (°C)	
					Linear mycelial growth	Mycelial density increase
<i>H18</i>	Hokkaido	43°24'N	142°36'E	6.2	20.6	17.8
<i>Rin10</i>	Hokkaido	43°22'N	143°59'E	3.9	17.1	17.3
<i>I122</i>	Iwate	39°56'N	141°14'E	9.4	20.8	17.7
<i>I129</i>	Iwate	39°56'N	141°14'E	9.4	20.3	17.3
<i>I33</i>	Iwate	39°56'N	141°14'E	9.4	20.5	17.7
<i>Sakai</i>	Kyoto	35°11'N	135°20'E	14.3	18.9	15.6
<i>NF2970</i>	Nara	34°10'N	135°41'E	14.6	19.6	19.8
mean±SD					19.6±1.3	17.6±1.2

<sup>(a)</sup>Normal annual mean air temperature observed at the nearest weather station of Japan meteorological agency from each study site



**Fig. 1. Flat-bottom vials containing the soil medium used in this study**  
The panel on the left shows the soil medium before inoculation and that on the right shows the medium after 89 day of cultivation at 20°C for strain I129

flasks containing the MYPG liquid medium and incubated at 23°C for 34 day under shaking at 35 rpm. Consequently, the mycelia were homogenized in a flask using Physcotoron homogenizer (Microtec Co. Ltd., Japan) at 1800 rpm for 5 s to prepare a uniform mycelial suspension for the next inoculation and incubated under static conditions continuously in the same flask. After 23 day of culture, 2 mL of the liquid medium including small pellets of *T. matsutake* mycelia was inoculated onto the soil medium, and the vial was capped and sealed with Parafilm (Bemis Flexible Packaging, USA). The cultures were incubated at 5, 10, 15, 20, 25, and 30°C for 89 day. Five replicates were kept for each strain at each temperature. All the incubations were carried out in the dark and under aseptic conditions.

### 2.3 Measurement of Mycelial Growth and Biomass

After 89 d of cultivation, linear growth of the *T. matsutake* mycelia from the inoculum was measured on four vertical lines under a dissecting microscope. The observed values were averaged and divided by the cultivation day number to obtain the growth rate for each vial. Thereafter, the soil medium containing the inoculated and/or cultured mycelia was collected immediately and stored at -30°C for further qPCR analysis. *T. matsutake* mycelial densities in the soil medium in all the incubation vials were analyzed by qPCR to determine the mycelial biomass [17]. The soil samples were lyophilized, pulverized with a Multi-beads Shocker (Yasui Kikai Corporation, Japan), and the sample DNA was extracted using the cetyltrimethylammonium-bromide (CTAB) lysis buffer. qPCR was performed using a LightCycler System (Roche Life Science, Germany) and the primer set, MY201f: 5'-GAGACACAACGGCGAGATT-3' and MY101r: 5'-ACCCTTACCCGCTCAGT-3', to amplify a 202-bp DNA fragment used to specifically quantify the *T. matsutake* mycelia [17]. DNA amplification and detection were performed in glass capillaries in a total volume of 20 µL, which contained 2 µL of LightCycler FastStart DNA Master SYBR Green I (Roche Diagnostics, Switzerland). The reaction mixture contained 3.2 µL (5 mM) of MgCl<sub>2</sub>, 1 µL (0.25 µM) of each primer, 10.8 µL of H<sub>2</sub>O (sterile PCR grade), and 2 µL of template or sample DNA. To detect the signal derived from *T. matsutake* mycelia, qPCR was performed using the following program, as recommended by the manufacturer: 1 cycle at

95°C for 10 min, followed by 50 cycles at 95°C for 10 s, 67°C for 2s, 72°C for 1 s, and 83°C for 1s, followed by signal detection. Unless stated otherwise, the standard curve was constructed using known numbers of copies of the template plasmid containing the target DNA. The copy number in the serial dilutions of the template plasmid ranged from 10<sup>7</sup> to 10<sup>1</sup>. The sample DNA was serially diluted and added to the 20-µL reaction mixture. The PCR products with a single melting curve that fitted the respective standard curve were considered as authentic, quantified PCR products.

### 2.4 Data Analysis


The mycelial growth rate and density in each vial were fitted to quadratic and tertiary equation models to estimate the optimum temperature for the mycelial growth or increase in the density of *T. matsutake*, because the observed value of mycelial growth rate and density showed a single peak with a long tail for some strains (Figs. 2 and 3). Next, the models were compared using the Akaike information criterion (AIC) (1) to select the most rational model for each strain:

$$\text{AIC} = \text{residual deviance} + 2(\text{the number of variances}) \quad (1)$$

Finally, the selected models were differentiated to estimate the optimum temperature. All the above-mentioned analyses were performed using R 2.14.0 [18].


## 3. RESULTS AND DISCUSSION

For all the *T. matsutake* strains tested, a linear mycelial growth was observed at 5, 10, 15, 20, and 25°C on the soil medium, but not at 30°C (Fig. 2). The mean linear mycelial growth rate was highest at 20°C for six of the seven strains. The daily linear growth rate at 20°C varied from 0.12 mm/d (strain Rin10 and I33) to 0.19 mm/d (strain I129). Significant difference among the strains was observed for the mean daily linear growth rate at 20°C ( $p = .01$ , ANOVA). The linear growth rate declined rapidly between 25°C and 30°C, which is in accordance with the results of a previous report [11]. The results of curve fitting showed that AIC was the smallest in the tertiary equation model for all the seven strains (Fig. 2). Using the tertiary equation model, the optimum temperature for linear mycelial growth was estimated to range from 17.1°C (strain Rin10) to 20.8°C (strain I122), with an average of 19.6 ± 1.3 SD°C (Table 1).



**Fig. 2. Effect of temperature on linear mycelial growth of seven *Tricholoma matsutake* strains in the soil medium after 89 d of cultivation**

Circles indicate the observed values, and lines indicate the approximate curve. The fitted model was selected from the quadratic or tertiary equation by comparing the akaike information criterion (AIC) for each strain (underlined)



**Fig. 3. Effect of temperature on mycelial density increase in seven *Tricholoma matsutake* strains in the soil medium after 89 d of cultivation**

Circles indicate the observed values, and lines indicate the approximate curve. The fitted model was selected from the quadratic or tertiary equation by comparing the akaike information criterion (AIC) for each strain (underlined)

The mycelial density increased during the incubation period at each selected temperature (Fig. 3). Consistent with the linear mycelial growth rate, significant difference among strains was observed for the mycelial growth rate at 20°C ( $p = .05$ , ANOVA). A decrease in mycelial density was observed between 25°C and 30°C as was observed for the linear mycelial growth rate. The curve fitting revealed the smallest AIC for the tertiary equation model for strain H18 and strain NF2970, and for the quadratic equation model for the other strains (Fig. 3). Thus, using a suitable growth model for each strain, the optimum temperature for mycelial density increase was estimated to range from 15.6°C (for the strain Sakai) to 19.8°C (strain NF2970), with the average being  $17.6 \pm 1.2^\circ\text{C}$  (Table 1). The mean optimum temperature for the mycelial density increase was slightly lower than that required for linear mycelial growth ( $p = .01$ ,  $t$ -test). No significant relationship was observed between the optimum temperature for mycelial density increase and that for linear mycelial growth ( $r = 0.24$ ,  $p = .05$ ). Although the linear mycelial growth was not observed at 30°C, the mycelial density increased at 30°C; this may be attributed to the inclusion of the mycelia present in the inoculum in the measurement of mycelial biomass. However, the reason for the difference between optimum temperatures for linear mycelial growth and mycelial density increase is unclear; the difference in the mechanism underlying linear mycelial growth and mycelial density increase probably affects the response to the temperature.

No clear relationship was observed between the optimum temperature for linear mycelial growth or mycelial density increase and the mean air temperature at the collection site ( $r < 0.2$ ,  $p = .05$ ), suggesting that the growth rate of strains collected from the cold region might not always be lower than that of strains collected from the warm region, as reported previously [19]. On the other hand, the distribution area of *T. matsutake* is known to be remarkably larger compared to the area included for strain collection in this study, such as that from the northern Europe [4], with an annual mean temperature of 4.4–6.7°C in Finland [20] to south-east Asia [3], with an annual mean temperature of 19.6°C in Xien Khouang city, Laos [21]. The inclusion of a wider area may provide greater variance of ecological conditions, especially temperature, among the growth sites. Therefore, further studies comparing samples collected from a wider area are warranted to verify the relationship between

the temperature of the natural growth site and the optimum temperature for mycelial growth.

Thus, in this study, we cultivated the *T. matsutake* mycelia in a soil medium with conditions similar to those in the natural habitat and verified the optimum temperature for *T. matsutake* growth with respect to linear mycelial growth and mycelial density increase. The optimum temperatures for linear mycelial growth and mycelial density increase for each of the strains tested were about 3–5°C lower than those reported in previous studies on *T. matsutake* using agar media (such as 23°C) [11] or using liquid media (such as 25°C) [22]. The estimated optimum temperatures were also lower than those for most of the 62 species of ectomycorrhizal fungi grown in liquid medium [23] and for the two allied species of *T. matsutake*, namely *T. fulvocastaneum* [24] and *T. bakamatsutake* [25], grown on agar media. The optimum temperatures lower than those reported previously are consistent with the climatic conditions prevailing at the isolation sites because the ground temperature does not always reach 20°C, even in summers, especially in the northern mainland of Japan [26].

#### 4. CONCLUSION

The optimum temperatures for linear mycelial growth and mycelial density increase of *T. matsutake* were lower than those reported previously using agar or liquid media. The determination of the optimum temperatures would contribute to the improvement of artificial cultivation methods for *T. matsutake*. Based on the results of this study, the optimum temperature for cultivation of *T. matsutake* for use as an inoculum for forest or nursery stock is expected to be 18°C.

#### ACKNOWLEDGEMENTS

We thank Dr. Akira Ohta, Shiga prefecture, Japan, for technical advice with laboratory analyses. We are also grateful to Akiko Takou and Yuki Hirakawa for assistance with the experiments. This work was partly supported by a grant titled “Technology development for the optimal use of forest resources” from the Ministry of Agriculture, Forestry and Fisheries of Japan.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Imazeki R, Hongo T. Colored illustrations of mushrooms of Japan. In: Imazeki R, Hongo T, Editors. Colored Illustrations of Mushrooms of Japan. Osaka: Hoikuisha Publishing. 1987;1:77. Japanese
2. Ogawa M. Microbial ecology of mycorrhizal fungus, *Tricholoma matsutake* Ito et Imai (Sing.) in pine forest, 1: Fungal colony (Shiro) of *Tricholoma matsutake*. Bull Gov For Exp Stn. 1975;272:79–121. Japanese
3. Yamanaka K, Aimi T, Wan J, Cao H, Chen M. Species of host trees associated with *Tricholoma matsutake* and close allies in Asia. Mushroom Sci Biotechnol. 2011;19(2):79–87. Japanese
4. Vaario LM, Savonen EM, Peltoniemi M, Miyazawa T, Pulkkinen P, Sarjala T. Fruiting pattern of *Tricholoma matsutake* in Southern Finland. Scand J For Res. 2015;30(4):259–65.
5. Yamanaka T. Researches for development of the cultivation of 'matsutake', a prized mushroom produced by the ectomycorrhizal basidiomycete *Tricholoma matsutake*. Bull For For Prod Res Inst. 2012;11(3):85–95. Japanese
6. Wang Y, Ian RH. Edible ectomycorrhizal mushrooms: Challenges and achievements. Can J Bot. 2004;82(8):1063–73.
7. Narimatsu M, Koiwa T, Masaki T, Sakamoto Y, Ohmori H, Tawaraya K. Relationship between climate, expansion rate, and fruiting in fairy rings ('shiro') of an ectomycorrhizal fungus *Tricholoma matsutake* in a *Pinus densiflora* forest. Fungal Ecol. 2015;15:18–28.
8. Carlile M, Watkinson S, Gooday G. The fungi. Second. London: Academic Press; 2001.
9. De la Varga H, Águeda B, Ágreda T, Martínez-Peña F, Parladé J, Pera J. Seasonal dynamics of *Boletus edulis* and *Lactarius deliciosus* extraradical mycelium in pine forests of Central Spain. Mycorrhiza. 2013;23(5):391–402.
10. Ohta A. A new medium for mycelial growth of mycorrhizal fungi. Trans Mycol Soc Japan. 1990;31(3):323–34.
11. Hamada M. Matsutake. Shizen. 1953;8(10):56–64. Japanese
12. Yamada A, Kanekawa S, Ohmasa M. Ectomycorrhiza formation of *Tricholoma matsutake* on *Pinus densiflora*. Mycoscience. 1999;40:193–8.
13. Fukasawa Y. Temperature effects on hyphal growth of wood-decay basidiomycetes isolated from *Pinus densiflora* deadwood. Mycoscience. 2018;59(3):259–62.
14. Dowson CG, Boddy L, Rayner ADM. Development and extension of mycelial cords in soil at different temperatures and moisture contents. Mycol Res. 1989;92(4):383–91.
15. Sato T, Yaegashi K, Ishii S, Hirano T, Kajiwara S, Shishido K, et al. Transformation of the edible basidiomycete *Lentinus edodes* by restriction enzyme-mediated integration of plasmid DNA. Biosci Biotechnol Biochem. 1998;62(12):2346–50.
16. Yamada A, Katsuya K. Mycorrhizal association of isolates from sporocarps and ectomycorrhizas with *Pinus densiflora* seedlings. Mycoscience. 1995;36(3):315–23.
17. Yamaguchi M, Narimatsu M, Fujita T, Kawai M, Kobayashi H, Ohta A, et al. A qPCR assay that specifically quantifies *Tricholoma matsutake* biomass in natural soil. Mycorrhiza. 2016;26(8):847–61.
18. R Development Core Team. R: A language and environment for statistical computing; 2015.
19. Kawai M, Abe S. Studies on the artificial reproduction of *Tricholoma matsutake* (S. Ito et Imai) Sing. I. Effects of carbon and nitrogen sources in media on the vegetative growth of *T. matsutake*. Trans Mycol Soc Japan. 1976;17:159–67. Japanese
20. Vaario L, Yang X, Yamada A. Biogeography of the Japanese gourmet fungus, *Tricholoma matsutake*: A review of the distribution and functional ecology of matsutake. In: Tedersoo L, Editor. Biogeography of Mycorrhizal Symbiosis. Cham: Springer. 2017;319–44.
21. World Weather Online; 2019. (Accessed 5 November 2019) Available:<https://www.worldweatheronline.com/>
22. Lee W, Han S, Kim B, Shrestha B, Lee S, Ko C, et al. Proliferation of *Tricholoma matsutake* mycelial mats in pine forest using mass liquid inoculum. Mycobiology. 2007;35(2):54–61.
23. Akama K, Okabe H, Yamanaka T. Growth of ectomycorrhizal fungi on various culture media. Bull For For Prod Res Inst. 2008;7(4):165–81. Japanese

24. Todo C, Torigoe S. Studies on cultivation method of *Tricholoma fulvocastaneum* (I) The physiological characteristics. Bull Hyogo Prefect For Technol Cent. 1996;49:1–4. Japanese
25. Jeon S-M, Ka K-H, Hong K-S. Mycelial growth and *in vitro* ectomycorrhizal synthesis on *Pinus densiflora* seedlings of *Tricholoma bakamatsutake* in Korea. Korean J Mycol. 2014;42(4):312–21.
26. Omori H. The affect of weather conditions on the production volume of matsutake in Iwate prefecture. Bull Iwate Prefect For Technol Cent. 1997;7:67–72. Japanese

© 2019 Narimatsu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<http://www.sdiarticle4.com/review-history/52395>